



NATURE

A Teeny Tiny Amphibian Had The Earliest Known Example of a Rapid-Fire Tongue

like amphibians you've likely never heard of.

Now extinct, albies had a dream run. They'd been around since the Middle Jurassic around 165 million years ago, and probably even earlier. They lived through the age of [dinosaurs](#) (and saw out their extinction), then lived through the rise of the great apes, before quietly disappearing about 2.5 million years ago.

they lived.

[New research](#) by my colleagues and I, published today [November 5] in *Science*, reveals these amphibians were the earliest known creatures to have rapid-fire tongues. This also helps explain why albies were once misidentified as chameleons.

A miniature marvel uncovered

The reason albies remained largely elusive until recently is because they were tiny. Their slight, fragile bones are usually found as isolated jaw and skull fragments, making them hard to study.

The first almost complete albie specimen was found in the wetland environment deposits of Las Hoyas, Spain, and [reported in 1995](#). Even though it was squashed flat, it was enough for palaeontologists to conclude albies were unlike any living salamander or any other amphibian.

They were completely covered in scales like reptiles, had highly flexible necks like mammals, an unusual jaw joint, and large eye sockets suggesting good vision. Why were albies so unique?

Mistakes do happen

The answer partly came to light in 2016, when a group of researchers [published a paper](#) demonstrating the diversity of lizards found in the

skin, claws, and muscles, still attached within the fossilised tree resin.

The researchers used "[micro-CT](#)" technology to digitally excavate and study the specimens in detail. This involved using 3D imaging to digitally remove the fossil from the amber and study it on a computer – a technique that avoids the risk of physically damaging the fossil.

They noticed one small, juvenile specimen had a long rod-shaped tongue bone. It was identified as the earliest known chameleon: a remarkable discovery! Or was it?

Alas, mistakes do happen in science. As lizard experts, the researchers had interpreted their results through this lens. It took the keen eye of Susan Evans, a professor of vertebrate morphology and palaeontology at University College London, to recognise this particular "lizard" was actually a misidentified albie.

A tongue-tying revelation

Some time later, Sam Houston State University assistant professor Juan Daza spotted another unbelievable specimen among a collection of fossils preserved in [Burmite amber](#), [ethically sourced](#) from Myanmar's Kachin state.

[Australian Synchrotron](#) in Melbourne.

Named after a class of mythical spirits responsible for guarding natural treasures, Yaksha, and the person who discovered the fossil, Adolf Peretti (founder of the non-profit [Peretti Museum Foundation](#)) — the *Yaksha perettii* specimen was an entire skull trapped in golden amber.

Quick hits to unsuspecting prey

Its features that stood out were a long bone projecting back out of the mouth and soft tissue remains, including part of the tongue, jaw muscles, and eyelids. By sheer luck, the soft tissue remains proved the long bone in the mouth was directly attached to the tongue.

In other words, *Y. perettii* was a predator armed with an incredible weapon: a specialised ballistic tongue that fired at [lightning](#) speed to capture prey – just as chameleons do today. It's no wonder the original juvenile, only 1.5 centimetres (0.59 inches) long, was initially mistaken for a chameleon.

Chameleon Tongue In Slow Motion | One Life | BBC Earth



We believe albies' projectile tongues were just as fast, used to great effect while sitting motionless in trees or on the ground. If so, this also explains why albies had unusual jaw joints, flexible necks, and large, forward-facing eyes. All these traits would have made up their predator toolkit.

Tree sap turned to iridescent amber

Despite these remarkable new insights, however, many mysteries of albanerpetontids remain. For instance, how exactly are they related to other amphibians? How did they survive for so long, only to die out relatively recently?

We'll need more intact specimens to answer these questions. And most of these specimens will probably come from the [Hukawng Valley](#) in Kachin, Myanmar.

It's expected about 100 million years ago this region was an island covered in vast forests. Global temperatures back then would have exceeded today's, with trees producing vast amounts of resin (which later turned into amber) as a result of damage by insects and fire.

Amber studied from this region will not only increase our knowledge of its expired ecosystems, it could also provide insight into how certain organisms today might evolve in response to a warming climate.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).
